

Chapter 1: Multivariables Functions

- 1.1 Functions of Two Variables
- 1.1.1 Function representations
- 1.1.2 3-D Coordinate System
- 1.1.3 Graph of two variable functions
- 1.1.4 Sketching of the function (3-D

*Level Curves

- 1.1.5 Domain and Range
- 1.2 Functions of Three Variables
- 1.2.1 Domain and Range
- 1.2.2 Level Surfaces

1.1 Functions of Two Variables

$$z = f(x, y)$$

Means that z is a function of x and yin the sense that a unique value of the **dependent variable** z is determined by specifying values for the **independent variables** x and y.

> $(x, y) \in Domain$ $z \in Range$

and

x and y: the two different independent variables

z: the dependent variable

Domain (D) : the set of all possible inputs (x, y) of the function f(x, y) that is

Range (R) : the set of output z that result when (x, y) varies over the domain D

For example,

1.
$$f(x, y) = \sqrt{x^2 + y^2}$$

$$f(1,1) = \sqrt{1^2 + 1^2} = \sqrt{2}$$

Function of two variables

Substitute 1 for *x* and 1 for *y*

2.
$$z = f(x, y) = \sqrt{64 - x^2 + e^{xy}}$$

 $f(1,0) = \sqrt{64 - 1 + 1} = 8$
 $f(2,-3) = \sqrt{64 - 4 + e^{-6}} = \sqrt{60 + e^{-6}}$

1.1.1 Function Representation of z = f(x, y)

3-D coordinate system



f(x, y) is a rule that assigns a unique real number to each point (x, y) in same set D in the xy-plane

Coordinate Planes



1.1.2 3-D Coordinate system



3 DIMENSIONAL CARTESIAN COORDINATE SYSTEM

3D coordinate system has 3 main planes:-

xy plane	or	z = 0	(x, y, 0)
xz plane	or	y = 0	(x, 0, z)
yz plane	or	x = 0	(0, y, z)

The orientation of xyz-axis



1.1.3 Graph of a Function of Two Variables

The graph of the function f of two

variables is the set of all points (x, y, z) in three-dimensional space, where the values of (x, y) lie in the domain of *f* and z = f(x, y).





The graphs of z = f(x, y) is called a surface in 3D system or three-space (\Re^3).

It looks like a blanket!

Four types of surface in space:

1.1.3.1 Planes

Example 1 z = 0, y = 0, x = 0x = 3, y = -1, z = 5

Given as a constant equation with **onevariable.**

Example 2

y = -x + 6, 2y = 4z + 5, z + x = 4Given as a linear equation with **two-variable**.

Example 3 Tetrahedron

$$y + x + y = 1$$

z = 6 - 3y + 2x

Given as a linear equation with **three**-**variable**.



How to sketch of the given functions

- 1) Determine the variables
- 2) Sketch the trace in coordinate planes (based on the variables exist)
- 3) Make the projection onto the traceplane which is parallel to the (variables which is not exists)-axis





Eg 7: Sketch the graph of 2+2+4 = 1. Solution : the traces in the coordinate planes : -> yz-plane, x = 0 : the straight line 4=1-2. -> x 2 - plane, y = 0 : the straight line 2 = 1-x. -> xy-plane, z=0: the straight line -> 2+ 2+ 445=1 51

Exercise 1 : Sketch the graph of 2 : 6 - by + 2x



1.1.3.2 CURVED SURFACE (MONLINEAR BOUASION) (NONLINEAR BOUASION) (NONLINEAR BOUASION) $x^{2}+y^{2}=x^{2}$ $y^{2}-z^{2}=9$

the given eq is have two variables. firstly, sketch the graph trace in a plane (based on the given variables)



the given eq's have three variables firstly, sketch the graph traces in the three planes (coordinate planes)



Sketch the graph trace in a plane (based on the given Vanables)













14.4 Surfaces in Space 647



FI

.

 $(14)_{s}$

How to sketch curved surfaces ?

Level Curves

To sketch the graph of two variables, we need to familiar with the contour maps.

Notice that when the plane z = C intersects with the surface z = f(x, y), the result is the space curve with the equation $\frac{f(x,y)=C}{x}$, so we called these as the **level curves**.

1.1.4 Sketch of the surface z = f(x, y)

* the set of point (x, y) in xy-plane that satisfy f(x, y) is called **level curves** /contour curves





Sketching surfaces with level curves

Let z = f(x, y) is a function of two variables

- ✓ Plane z = C intersects with the surface $z = f(x, y) \rightarrow f(x, y) = C$
- ✓ The set of point (x, y) in the xy-plane that satisfy f(x, y) = C is called the level curve of f at C
- ✓ An entire family of level curves is generated as *C* varies over the range of *f* ✓ The graph of *z* = *f*(*x*, *y*) is a surface which can be obtained by sketching the contour map (set of level curves) on xyplane

Example

Sketch the contour lines/level curves and the graphs

(i) $z = x^2 + y^2$, c = 0, 1, 2, 3, 4, 9

(ii)
$$z = \sqrt{x^2 + y^2}$$
, $c = 0, 1, 4, 9$

(iii)
$$z = 6 - x^2 - y, c = 0, 2, 4, 6$$

Solution

(i)
$$z = x^2 + y^2$$
, $c = 0, 1, 2, 3, 4, 9$

Sketching the level curves

- first, replace z with the value of c

- second, plot the graph on the xy-plane



The traces in the coordinate planes:

- *yz*-plane, x = 0: the quadratic curve, $z = y^2$
- *xz*-plane, y = 0: the quadratic curve, $z = x^2$

• *xy*-plane,
$$z = 0$$
: a point (the origin)





(ii)
$$z = \sqrt{x^2 + y^2}$$
, $c = 0, 1, 4, 9$

c = 0	$:\sqrt{x^2+y^2}=0$	level curves
<i>c</i> =1	$: \sqrt{x^2 + y^2} = 1$	in xy-plone
<i>c</i> =4	$: \sqrt{x^2 + y^2} = 4$	×
<i>c</i> =9	$: \sqrt{x^2 + y^2} = 9$	

The traces in the coordinate planes:

- *yz*-plane, x = 0: the straight line, z = y
- *xz*-plane, y = 0: the straight line, z = x
- *xy*-plane, z = 0: a point (the origin)
- parallel to *xy*-plane, z = 4: the circle $x^2 + y^2 = 4^2$





eg.
$$Z = \int x^2 + y^2$$

the graph of
the E^2 given
(surface) and
(surface) and
 $I = \int x^2 + y^2$
 $Z = \int x^2 + y^2$
 $Z = 1$
 $Z = 1$

(ii) $z = 6 - x^2 - y$, c = 0, 2, 4, 6.

Sketching the level curves

- first, replace z with the value of c
- second, plot the graph on the xy-plane

c = 0	$: 6 - x^2 - y = 0 \Longrightarrow y = -x^2 + 6$
<i>c</i> =2	$: 6 - x^2 - y = 2 \Longrightarrow y = -x^2 + 4$
<i>c</i> =4	$: 6 - x^2 - y = 4 \Longrightarrow y = -x^2 + 2$
<i>c</i> =6	$: 6 - x^2 - y = 6 \Longrightarrow y = -x^2$





1.1.5 Domain and Range of z = f(x, y)

Domain :
$$\{(x, y) | x \in \mathbb{R}, y \in \mathbb{R}, \underline{???}\}$$

 \uparrow
any constraint

??? f(x, y) may consist:



*Sometimes we need to sketch the domain of the function given.

Range – z-values that results when (x,y) varies over the domain

- (i) z positive ?
- (ii) z negative ?
- (iii) z zero?
- (iv) z has maximum value ?
- (v) z has minimum value ?

Range : $\{z \mid z \in \mathbb{R}, \underline{???}\}$

put the limitation of z here!!

Example

Describe the domain and the range of $z = \sqrt{64 - 4x^2 - y^2}$.

Solution



Domain: $\{(x,y) \mid x \in \mathbb{R}, y \in \mathbb{R}, Gu - ux, y^{2} = 0\}$ or Jomain: $\{(x,y) \mid x \in \mathbb{R}, y \in \mathbb{R}, \frac{x}{16} + \frac{y}{16} \leq 1\}$ Ronge: $\{z \mid z \in \mathbb{R}, 0 \leq z \leq 5 \leq 164\}$ ov Ronge: $\{z \mid z \in \mathbb{R}, 0 \leq z \leq 8\}$

Example

Find the domain and range of $z = x^2 \sqrt{y-1}$. Solution Construint : 430 Domain : [(x.y) x EIR, Y EIR, 4201 Ronge : xJy > 0 (always tve) 23-1 => Ronge : {2 | 2 EIR, 2 3 - 1 3 Sketching of domain

Example

Find the domain and the range of $z = \ln(x^2 - y)$. Solution



Example Find the domain and the range of $z = 4 - x^2 - y^2$.

Solution

Domain : $\{(x, y) | x \in \mathbb{R}, y \in \mathbb{R}\}$

Range: $\{z \mid z \in \mathbb{R}, z \le 4\}$



1.2 Functions of Three Variables 1.2.1 Domain and Range

Definition

A function *f* of three variables is a rule that assigns to each ordered triple (x, y, z)in some domain *D* in space a unique real number w = f(x, y, z).

The range consists of the output values for w.

Example 1

Identify the domain and range for the following functions.

a).
$$w = \sqrt{x^2 + y^2 + z^2}$$

 $x^2 + y^2 + z^2 \ge 0$ for all points in space.
Domain : entire space

Domain : { $(x, y, z) | x \in \mathbb{R}, y \in \mathbb{R}, z \in \mathbb{R}, x^2 + y^2 + z^2 \ge 0$ }

Range : $[0, \infty)$ Range : $\{w | w \in \mathbb{R}, w \ge 0\}$

b)
$$w = \sqrt{1 - (x^2 + y^2 + z^2)}$$

We must have $1 - (x^2 + y^2 + z^2) \ge 0$ in order to have a real value for f(x, y, z). Rewriting the condition, we obtained

$$x^2 + y^2 + z^2 \le 1$$

Thus the domain consists of all points on or within the sphere $x^2 + y^2 + z^2 = 1$, or Domain : $\{(x, y, z) | x \in \mathbb{R}, y \in \mathbb{R}, z \in \mathbb{R}, x^2 + y^2 + z^2 \le 1\}$ Range : [0, 1] or Range : $\{w | w \in \mathbb{R}, 0 \le w \le 1\}$

c)
$$w = \frac{1}{x^2 + y^2 + z^2}$$

Domain : $\{(x, y, z) : (x, y, z) \neq (0, 0, 0)\}$ or

Domain : { $(x, y, z) | x \in \mathbb{R}, y \in \mathbb{R}, z \in \mathbb{R}, x^2 + y^2 + z^2 \neq 0$ }

Range : $(0, \infty)$ or

Range : $\{w | w \in \mathbb{R}, w > 0\}$

d) $w = xy \ln z$ Domain : $\{(x, y, z) | x \in \mathbb{R}, y \in \mathbb{R}, z \in \mathbb{R}, z > 0\}$ Range : $(-\infty, \infty)$ or Range : $\{w | w \in \mathbb{R}, -\infty \le w \le \infty\}$

1.2.2 Level Surfaces

The graphs of **functions of three variables** consist of points (x, y, z, f(x, y, z)) lying in **four-dimensional space**.

- Graphs cannot be sketch effectively in three-dimensional frame of reference.
- Can obtain insight of how function behaves by looking at its threedimensional level surfaces.

The graph of the equation f(x, y, z) = kwill generally be a **surface in 3-space** which we call the **level surface** with constant *k*.

Remark

The term "level surface" is standard. It need **not** be level in the sense being horizontal; it is simply a surface on which all values of *f* are the same.

Example

Describe the level surfaces of (a) $f(x, y, z) = x^2 + y^2 + z^2$ (b) $f(x, y, z) = z^2 - x^2 - y^2$

Solution

(a)
$$f(x, y, z) = x^2 + y^2 + z^2$$

The level surfaces have equation of the form

$$x^2 + y^2 + z^2 = k$$

For k > 0, the graph of this equation is a sphere of radius \sqrt{k} , centred at the origin.

For k = 0, the graph is the single point (0, 0, 0).





Level surfaces of $f(x, y, z) = x^2 + y^2 + z^2$

b)
$$f(x, y, z) = z^2 - x^2 - y^2$$

The level surface have equation of the form

$$z^2 - x^2 - y^2 = k$$

For k > 0, the graph is a hyperboloid of two sheets.

For k = 0, the graph is a **cone**.

For k < 0, the graph is a hyperboloid of one sheet.

Level surfaces of



Exersizes

Describe the level surfaces of (i) $f(x, y, z) = x^2 + y^2$ for C = 4, C = 9.

(ii) $f(x, y, z) = 4x^2 + y^2 + 4z^2$ for w = 1, w = 4.