## **CHAPTER 2: Infinite Series**

- 2.1 Definition of Infinite Series
- 2.2 Telescoping and Geometric Series
- 2.3 The Integral Test
- 2.4 Divergence Test
- 2.5 Comparison Test
- 2.6 Root and Ratio Tests
- 2.7 Alternating Series Test
- 2.8 Absolute Convergence

### 2.1 Definition of Infinite Series

$$\{a_1, a_2, a_3, \cdots, a_n, a_{n+1}, \cdots\} \text{ - infinite sequence}$$

$$S_1 = a_1$$

$$S_2 = a_1 + a_2$$

$$S_3 = a_1 + a_2 + a_3$$

$$\vdots$$

$$S_k = a_1 + a_2 + a_3 + \cdots + a_k = \sum_{n=1}^k a_n$$
The sequence of partial sum,  $\{S_k\} = \text{Infinite}$ 
series or for short series only.

If there is a real no. S such that 
$$\lim_{k\to\infty} S_k = S$$
,  
that is  $\sum_{n=1}^{\infty} a_n = S$ .  
= the sum of  
the series

Then we say that the series converges,

$$\sum_{n=1}^{\infty} a_n \text{ converges.}$$

$$\bigotimes_{k\to\infty} If \lim_{k\to\infty} S_k \text{ does not exist or } \lim_{k\to\infty} S_k = \pm\infty,$$
Then the series 
$$\sum_{n=1}^{\infty} a_n \text{ diverges.}$$

### 2.2 Telescoping and Geometric Series

#### 2.2.1 Telescoping Series

A series  $\sum_{n=1}^{\infty} a_n$  is a telescoping series if there is a sequence  $\{b_n\}$  such that

$$a_n = b_n - b_{n+1}$$
;  $n = 1, 2, 3, \dots$ 

Then  $\sum_{n=1}^{\infty} a_n = b_1 - \lim_{n \to \infty} b_n$ . Hence the telescoping series converges if and only if the sequence  $\{b_n\}$  converges.

# Example

Show that the following series is a telescoping series. Hence, determine the series converges or diverges.

(a) 
$$\sum_{n=1}^{\infty} \frac{1}{n \quad n+1}$$
  
(b)  $\sum_{n=1}^{\infty} \ln\left(\frac{n+2}{n+3}\right)$ 

# Example



$$\sum_{n=0}^{\infty} ar^{n} = a + ar + ar^{2} + ar^{3} + \dots ; \quad a \neq 0$$

Note: *a* is the first term.

The geometric series is convergent if |r| < 1 and its sum is  $\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r}$ . If |r| > 1, then the

geometric series is divergent.

# Example

(a) Find the sum of geometric series

$$5 - \frac{10}{3} + \frac{20}{9} - \frac{40}{27} + \dots$$

(b) Is the series  $\sum_{n=0}^\infty 2^{2n} 3^{1-n}$  convergent or

divergent?

#### 2.3 The Integral Test

Let  $\sum_{n=1}^{\infty} a_n$  be a series with  $a_n > 0, n = 1, 2, ...$  and f

be a function such that  $f(n) = a_n$ . f is

continuous and decreasing function for all real  $x \ge 1$  and *L* is a real number.

(1) If 
$$\int_{1}^{\infty} f(x) dx = L$$
 then  $\sum_{n=1}^{\infty} a_n$  converges.  
(2) If  $\int_{1}^{\infty} f(x) dx = \infty$  then  $\sum_{n=1}^{\infty} a_n$  diverges.

Notes: \*This test cannot be used to calculate

sum. \*Use this test when f(x) is easy to

integrate. \*This test only applies to series that

have positive terms.

## Example

Determine whether the following series converges or diverges.

(a) 
$$\sum_{n=1}^{\infty} rac{1}{4n+5}$$
  
(b)  $\sum_{n=1}^{\infty} rac{1}{n^2}$ 

**Note:** The series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  is called *p*-series. The *p*-series converges if p > 1 and diverges if 0 .

## 2.4 Divergence Test

If 
$$\lim_{n\to\infty} a_n \neq 0$$
, then  $\sum_{n=1}^{\infty} a_n$  diverges. If  $\lim_{n\to\infty} a_n = 0$ ,  
then  $\sum_{n=1}^{\infty} a_n$  may be convergent or may be  
divergent, hence other tests should be used.  
Note: This test only determines the divergence  
of a series.

# Example

Show that the series diverges

(a) 
$$\sum_{n=1}^{\infty} \frac{2n}{n+1}$$
  
(b)  $\sum_{n=1}^{\infty} \cos n\pi$ 

(c) 
$$\sum_{n=1}^{\infty} \frac{e^n}{n}$$
  
(d)  $\sum_{n=1}^{\infty} \frac{n^{n+1/n}}{n+1/n}$ 

# 2.5 Comparison Test

Let 
$$\sum_{n=1}^{\infty} a_n$$
 and  $\sum_{n=1}^{\infty} b_n$  be series with nonnegative  
terms such that  
 $a_1 \leq b_1, a_2 \leq b_2, a_3 \leq b_3, \dots, a_n \leq b_n, \dots$   
for all  $n$ . If  $\sum_{n=1}^{\infty} b_n$  converges, then  $\sum_{n=1}^{\infty} a_n$   
converges. If  $\sum_{n=1}^{\infty} a_n$  diverges, then  $\sum_{n=1}^{\infty} b_n$  diverges.

### Notes:

\*The series that we use for comparison are usually the geometric series or the p-series. \*Use this test as a last resort; other tests are often easier to apply.

\*This test only applies to series with nonnegative terms.

# Example

(a) 
$$\sum_{n=1}^{\infty} \frac{n}{n^2 - \cos^2 n}$$
  
(b)  $\sum_{n=1}^{\infty} \frac{n^2 + 2}{n^4 + 5}$ 

Let  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  be series with positive terms

such that  $c = \lim_{n \to \pm \infty} \frac{a_n}{b_n}$ . If  $0 < c < \infty$ , then both

series converge or both diverge.

Note: This is easier to apply than the comparison test, but still requires some skill in

choosing the series  $\sum_{n=1}^{\infty} b_n$  for comparison.

If c = 0 and  $\sum_{n=1}^{\infty} b_n$  converges, then  $\sum_{n=1}^{\infty} a_n$  converges. If  $c = \infty$  and  $\sum_{n=1}^{\infty} b_n$  diverges, then  $\sum_{n=1}^{\infty} a_n$  diverges.

# Example

(a) 
$$\sum_{n=0}^{\infty} \frac{1}{3^n - n}$$
  
(b)  $\sum_{n=2}^{\infty} \frac{4n^2 + n}{\sqrt[3]{n^7 + n^3}}$ 

## **CHAPTER 2: Infinite Series**

- 2.1 Definition of Infinite Series
- 2.2 Telescoping and Geometric Series
- 2.3 The Integral Test
- 2.4 Divergence Test
- 2.5 Comparison Test
- 2.6 Root and Ratio Tests
- 2.7 Alternating Series Test
- 2.8 Absolute Convergence

### 2.6 Root and Ratio Tests

### 2.6.1 Root Test

Let  $\sum_{n=1}^{\infty} a_n$  be a series with positive terms such that  $R = \lim_{n \to \infty} \sqrt[n]{a_n}$ .

- (a) Series converges if R < 1
- (b) Series diverges if R > 1 or  $R = \infty$
- (c) No conclusion if R = 1 (Try another tests)

Note: Try this test when  $a_n$  involves n<sup>th</sup> powers.

# Example

(a) 
$$\sum_{n=1}^{\infty} \left( n^{1\!\!/n} - 1 
ight)^n$$

(b) 
$$\sum_{n=1}^{\infty} \frac{n^n}{3^{1+2n}}$$
  
(c)  $\sum_{n=1}^{\infty} \frac{n^2}{2^n}$ 

### 2.6.2 Ratio Test

Let  $\sum_{n=1}^{\infty} a_n$  be a series with positive terms and

suppose 
$$L = \lim_{n \to \infty} \frac{a_{n+1}}{a_n}$$

- (d) Series converges if L < 1
- (e) Series diverges if L > 1 or  $L = \infty$
- (f) No conclusion if L = 1 (Try another tests)

Note: Try this test when  $a_n$  involves factorials or n<sup>th</sup> powers.

# Example

Determine whether each series converges or diverges.

(a) 
$$\sum_{n=1}^{\infty} \frac{2^n}{n!}$$
  
(b)  $\sum_{n=1}^{\infty} \frac{2^n n!}{n^n}$ 

# 2.7 Alternating Series Test

Alternating series is a series of the form

$$\sum_{n=1}^{\infty} \ -1 \sum_{n=1}^{n-1} a_n = a_1 - a_2 + a_3 - a_4 + \dots$$

where  $a_n > 0$  for all n = 1, 2, 3, ...

The series converge if

(a) 
$$a_1 > a_2 > a_3 > a_4 > \dots$$

(b)  $\lim_{n\to\infty} a_n = 0$ 

Note: This test applies only to alternating

series. It is assumed that  $a_n > 0$  for all

 $n = 1, 2, 3, \dots$ 

### Example

(a) 
$$\sum_{n=1}^{\infty} rac{-1}{n}^{n+1}$$

(b) 
$$\sum_{n=1}^{\infty} \frac{-1^n n^2}{n^2 + 5}$$

(c) 
$$\sum_{n=1}^{\infty} \frac{\cos n\pi}{\sqrt{n}}$$

### 2.8 Absolute Convergence

#### Definition:

A series  $\sum_{n=1}^{\infty} a_n$  is called absolutely convergence if the series  $\sum_{n=1}^{\infty} |a_n|$  converges. A series  $\sum_{n=1}^{\infty} a_n$  is called conditionally convergence if the series  $\sum_{n=1}^{\infty} a_n$  converges, but the series  $\sum_{n=1}^{\infty} |a_n|$  diverges.

### Example

(a) 
$$\sum_{n=1}^{\infty} -1^{n-1} \frac{1}{\sqrt{2n+3}}$$
  
(b)  $\sum_{n=1}^{\infty} -1^n \frac{7}{n^3+1}$